

Effect of VASCEPA on Time to First Occurrence of Cardiovascular Events in Patients with Elevated Triglyceride levels and Other Risk Factors for Cardiovascular Disease in REDUCE-IT

	VASCEPA		Placebo		VASCEPA vs Placebo
	N = 4089 n (%)	Incidence Rate (per 100 patient years)	N = 4090 n (%)	Incidence Rate (per 100 patient years)	Hazard Ratio (95% CI)
Primary composite endpoint					
Cardiovascular death, myocardial infarction, stroke, coronary revascularization, hospitalization for unstable angina (5-point MACE)	705 (17.2)	4.3	901 (22.0)	5.7	0.75 (0.68, 0.83)
Key secondary composite endpoint					
Cardiovascular death, myocardial infarction, stroke (3-point MACE)	459 (11.2)	2.7	606 (14.8)	3.7	0.74 (0.65, 0.83)
Other secondary endpoints					
Fatal or non-fatal myocardial infarction	250 (6.1)	1.5	355 (8.7)	2.1	0.69 (0.58, 0.81)
Emergent or urgent coronary revascularization	216 (5.3)	1.3	321 (7.8)	1.9	0.65 (0.55, 0.78)
Cardiovascular death ^[1]	174 (4.3)	1.0	213 (5.2)	1.2	0.80 (0.66, 0.98)
Hospitalization for unstable angina ^[2]	108 (2.6)	0.6	157 (3.8)	0.9	0.68 (0.53, 0.87)
Fatal or non-fatal stroke	98 (2.4)	0.6	134 (3.3)	0.8	0.72 (0.55, 0.93)
[1] Includes adjudicated cardiovascular deaths and deaths of undetermined causality.					
[2] Determined to be caused by myocardial ischemia by invasive/non-invasive testing and requiring emergent hospitalization.					

FULL VASCEPA PRESCRIBING INFORMATION CAN BE FOUND AT WWW.VASCEPA.COM.